

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61L 25/00, A61K 38/36, C07K 14/75</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/33533</b> <b>(43) International Publication Date:</b> 6 August 1998 (06.08.98)
<b>(21) International Application Number:</b> PCT/EP98/00502 <b>(22) International Filing Date:</b> 30 January 1998 (30.01.98) <b>(30) Priority Data:</b> 97101569.8 31 January 1997 (31.01.97) EP <b>(34) Countries for which the regional or international application was filed:</b> DE et al. <b>(71) Applicant (for all designated States except US):</b> OMRIX BIOPHARMACEUTICALS S.A. [BE/BE]; 140, avenue Louise, B2, B-1050 Brussels (BE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> NUR, Israel [IL/IL]; Omri Laboratories Ltd., P.O. Box 619, 76106 Rehovot (IL). BAR, Lilliana [IL/IL]; Omri Laboratories Ltd., P.O. Box 619, 76106 Rehovot (IL). LIEBERMAN, Oded [IL/BE]; Omrix Biopharmaceuticals S.A., 140, avenue Louise, B2, B-1050 Brussels (BE). <b>(74) Agents:</b> MEYERS, Hans-Wilhelm et al.; P.O. Box 10 22 41, D-50462 Cologne (DE).	<b>(81) Designated States:</b> AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, DE, EE, GE, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> A STABILIZED MIXTURE COMPRISING FIBRINOGEN		
<b>(57) Abstract</b>  A stabilized solution of fibrinogen containing samples, in particular a stabilized solution of the biological active component (BAC) which is a solution of proteins derived from blood plasma comprising fibrinogen, tranexamic acid and arginine or lysine or mixtures of arginine and lysine, their pharmaceutically acceptable salts, as well as, optionally, substances forming a buffered solution in aqueous medium.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

A stabilized mixture comprising fibrinogen

The invention pertains to a mixture comprising fibrinogen, particularly to a stabilized solution of Biological Active Component (BAC), a two component tissue glue comprising separately components A and B as well as a fibrin clot obtainable by mixing component A and component B.

Fibrinogen containing samples can be used for various applications, e. g. in two component tissue glues or fibrin glues. Biologically Active Component (BAC) is a viscous solution of proteins containing fibrinogen typically in amounts about 50 mg fibrinogen per ml, which has been derived from cryoprecipitation of plasma. Such BAC is described in WO 94/22503. Since procedures employed for virus inactivation and concentration result in an increase of proteolytic activity in the cryoprecipitate, stabilization of the final product with anti-proteolytic agents is required. Some of these proteolytic enzymes are in a zymogen (pro-enzyme) form and their activation is promoted by the minute amount of activated enzymes present in the cryoprecipitate. The temperature range employed during separation of the cryoprecipitate has been found to enhance the activation of precursors of proteolytic enzymes. This activation presumably

takes place by a "cascade" sequence, whereby proteases activate the precursor forms of other proteolytic enzymes, including the generation of fibrinolytic plasmin from the precursor plasminogen. Inhibition of fibrinolysis and other proteolytic activity has been found to reduce the degradation of factor VIII and of other coagulative and adhesive proteins. It may occur that BAC coagulates spontaneously already after a storage for one day at 2 - 6 °C. At lower temperatures this process is slower; following storage for several months at -18°C, BAC without further treatment forms a solid clot after thawing and cannot be reconstituted. The same phenomenon has been observed with unpurified cryoprecipitate.

During the production process, plasminogen and other vitamin K dependent proteases are removed by aluminum hydroxide adsorption. However, some proteases are left in the final product.

It is desirable to stabilize fibrinogen comprised in a mixture, in particular a solution of BAC in order to ensure a save usage of the product when applied in clinical operations.

Surprisingly, this object is attained by a mixture comprising fibrinogen, 4-(aminomethyl)cyclohexane-carboxylic acid as well as arginine, lysine or combinations thereof. Preferably, the mixture is present as a solution of fibrinogen, tranexamic acid (4-(aminomethyl)-cyclohexane-carboxylic acid) as well as arginine, lysine or combinations thereof. In a preferred embodiment the invention comprises a solution of the biological active component (BAC) which has been stabilized with a combination of tranexamic acid its physiologically acceptable salts and arginine or lysine or combinations of lysine and arginine. Optionally, the solution is buffered to a physiological compatible pH value. Tranexamic acid is a protease inhibitor which scientific name

is 4-(aminomethyl)cyclohexane-carboxylic acid. According to the invention a buffer containing glycine is preferred.

Arginine as well as lysine can be employed according to the invention as a common salt, for example, as hydrochloride. The BAC is preferably obtained from concentrated cryoprecipitate after being worked up as disclosed in EP-A-534 178. Arginine has been described in the art as a stabilizer in therapeutic protein concentrates. However, the synergistic effect provided by the combination of tranexamic acid and arginine as well as lysine is surprising.

Preferably, the amount of tranexamic acid in the solution of BAC is from about 1 - 20% by weight. The amount of arginine or lysine hydrochloride is preferably from about 0.1 - 4% by weight. More preferred are amounts of tranexamic acid of from about 5 - 15% by weight, particularly preferred from about 8 - 12% by weight. Typically about 10% by weight can be used. The amount of arginine or lysine hydrochloride is more preferred in the range of from about 1 - 3% by weight, in particularly preferred are about 2% by weight.

The fibrinogen comprising sample, in particular a solution of BAC, is preferably derived from a cryoprecipitate which was concentrated by ultra filtration as described in WO-A-94/22503. The BAC preferably comprises a fibrinogen content of from about 15 - 150 mg/ml in particular of from 20 - 80 mg/ml. The amount of fibrinogen can be measured according to Clauss' Method (Clauss, A., "Gerinnungsphysiologische Schnellmethode zur Bestimmung des Fibrinogens", Acta. Haematol., 17, 237 - 247, 1957).

The use of a BAC derived from concentrated cryoprecipitate is advantageous since such a fraction contains besides fibrinogen also valuable blood components which play an important role for blood-clotting when a proteolytic enzyme such as human thrombin is contacted with a BAC solution.

Valuable components are factor VIII, factor XIII, fibronectin, vitronectin, von Willebrand factor (vWF), etc.

Preferably, the components are derived from cryoprecipitate, in particular concentrated cryoprecipitate. However, it is also possible that the components fibrinogen, factor VIII, factor XIII, fibronectin, von Willebrand factor (vWF), vitronectin have been prepared by recombinant methods. Such preparations, for example, for factor VIII are commercially available.

It was found that the combination of tranexamic acid and arginine or lysine stabilizes a mixture containing fibrinogen in particular a solution of the biological active component BAC, (at 2 - 6°C for at least 14 days by conserving at least 50 % of the fibrinogen activity). When only one of the components was used, the sample was significantly more instable.

Furthermore, it was surprising that the use of tranexamic acid and arginine or lysine hydrochloride did not adversely effect properties of a blood clot obtained from BAC stabilized according to the invention. For example, the maximum elongation of the clot was maintained also after 14 days of storage of the respective samples as well as the tensile force was almost maintained on the same level. Also the factor VIII activity in the BAC was not negatively affected when tranexamic acid and arginine or lysine were added.

In particular, the stabilized solution of the biological active component BAC according to the invention is well suitable for the preparation of a two component tissue glue. A tissue glue according to the invention is understood as a system which can be applied with or on a patient in need thereof, for example, to avoid severe bleeding during surgical operations. The tissue glue is also addressed as fibrin glue and was basically analogous to the natural blood

clotting cascade. The tissue glue is derived from two components prior to the application in surgical operations. One component contains fibrinogen which upon exposure to a proteolytic enzyme such as human thrombin forms fibrin which is the polymer forming the basic material of the natural blood clot. During surgical operations the two components are applied, for example, by two syringes which are emptied simultaneously by mixing the two components as fast as possible and avoiding the blockage of the supply lines. The solution according to the invention is advantageous for preparing a two component fibrin glue since the fibrinogen solution has not to be freshly prepared but can be stored in a refrigerator at  $-18^{\circ}\text{C}$ , neither losing its clotting ability nor its outstanding mechanical properties. The degrees of activity of fibrinogen can be balanced by providing a higher amount of fibrinogen in the solution so that the proper use of the fibrinogen containing solution (BAC) is not hampered.

Therefore, subject of the present invention is also a two component tissue glue comprising separately components A and B wherein component A comprises a solution according to the embodiments described in claims 1 to 10 and a component B comprising a solution of a proteolytic enzyme which is capable to react with fibrinogen (or BAC respectively) to form fibrin.

Preferably, the proteolytic enzyme is human thrombin in particular having an activity of from about 2 to 4,000 IU/ml. The activity of thrombin is measured according to the clotting assay (European Pharmacopoeia). It is understood by the skilled person that a fibrin glue may be defined by its content of clottable protein instead of the definition based on clottable fibrinogen.

In order to provide a balanced solution of the mixed components A and B it can be advantageous to add to component

B approximately the same concentration of tranexamic acid and arginine.

Preferably, the components A and B are applied in such a way that equal volumes of the two components are mixed and applied onto the patient at the side of the respective wound. Of course, it is to be understood that the two component tissue glue can be employed not only during surgical operations but also in other situations where bleeding must be stopped. The two components are preferably applied in a ratio of 1 : 1.

The following examples illustrate the advantages of the stabilizing combination of tranexamic acid and arginine. These examples are by no means limiting but explain the invention in greater detail.

#### Example 1

##### Preparation of test substance

To 800 ml of a sample 0.8 g of sodium azide (0.1%) will be added as a powder to the bulk to control bacterial growth, immediately after receiving the sample from the production line. The addition of a bacteriostat is necessary to prevent contamination since high concentration of fibrinogen in sample is difficult to filtrate.

Each concentration tested will be prepared extemporaneously by addition of a mixture of the two solid components to a 100 ml aliquot of the sample. The addition of tranexamic acid and arginine-HCl will be performed in a beaker with moderate stirring (multipoint magnetic stirring plate) for 10 minutes. All formulations tested will be prepared in parallel and their fibrinogen concentration will be adjusted to 50 mg/ml by addition of buffer B.



Group No.	BAC (ml)	Tranexamic Acid (g)	Arginine (g)	Buffer B (ml)	Final vol. (ml)
A	100	0	0	20	120
B	100	0	2.4	17.6	120
C	100	6	2.4	11.6	120
D	100	12	2.4	5.6	120
E	100	12	0	8	120
F	100	12	1.2	6.8	120
G	100	12	4.8	3.2	120

After an additional 5 minutes stirring, 10 ml siliconized glass vials will be filled with 5 ml aliquots. All vials will be frozen simultaneously and stored at -80°C until used. Before experiment, they will be placed at a temperature of 2 - 6°C (day "0", base-line time).

Two vials from each group will be kept at -80°C as positive controls. All experiments will be performed in duplicates and the vials will be labelled according to stabilizer concentrations as follows:

- GROUP A: no stabilizer-vials  $A_1$  -  $A_{20}$   
GROUP B: 0% tranexamic acid and 2% arginine monohydrochloride-vials  $B_1$  -  $B_{20}$   
GROUP C: 5% tranexamic acid and 2% arginine monohydrochloride-vials  $C_1$  -  $C_{20}$   
GROUP D: 10% tranexamic acid and 2% arginine monohydrochloride-vials  $D_1$  -  $D_{20}$   
GROUP E: 10% tranexamic acid and 0% arginine monohydrochloride-vials  $E_1$  -  $E_{20}$   
GROUP F: 10% tranexamic acid and 1% arginine monohydrochloride-vials  $F_1$  -  $F_{20}$   
GROUP G: 10% tranexamic acid and 4% arginine monohydro-

chloride-vials  $G_1$  -  $G_{20}$

Extra quantities of sample will be held as retention samples and used if needed.

Before testing all  $-80^{\circ}\text{C}$  frozen vials will be thawed at  $37^{\circ}\text{C}$  for 15 minutes and placed at  $2 - 6^{\circ}\text{C}$  (day "0"). Two vials from each group will be rolled at room temperature for 15 minutes at day 0 and their mechanical properties and biochemical tests will be performed and evaluated. The same procedure will be repeated on days 2, 4, 7, 9, 11, 14 and 30.

#### Mechanical Properties

Elongation test of fibrin glue will be done in a tensile machine CHATILLON MODEL TCD-200 with a CHATILLON Force Gauge of 1,000 g. Data will be handled by a computer program.

This instrument is a motor-driven tension and compression tester designed for testing the resilience, yield points and breaking strengths of various products and materials.

In order to produce standardized clots of solidified glue in a form which could easily be attached to the tension testing machine, a special cast was designed. This consists of two conical aluminum moulds, each of 2.5 cm in height, which are placed one on top of the other (see figure 1). The liquid components of the glue are injected into the cast where they solidify into a standard clot of 12.7 mm x 5 mm.

Fig.1 illustrates the special cast designed to hold the liquid glue until solidification .

The casts are then anchored to the tension testing machine, and the tensile strength and the elongation of the cylinder of glue within the cast measured by pulling the two casts apart.

A computer program monitors the tensile force and elongation

every 0.2 s. The points are plotted graphically on two axes: elongation versus force in grams at a given time.

#### Test procedure

Vials containing BAC were incubated at 37°C for 15 minutes and then rolled at room temperature for 15 minutes. Custom designed aluminum casts (as previously described) were filled with a combined solution of 0.5 ml of BAC and 0.5 ml of thrombin (8 IU/ml).

The casts were left at room temperature for 45 minutes to allow the glue to solidify, and then mounted onto the gauge and the resultant clots tested for maximum elongation capacity and tensile strength (tensile force required to break the clot).

#### Biochemical tests performed

The following proteins were tested for their stability by the methods described:

##### Fibrinogen clotting activity:

Fibrinogen was measured quantitatively by the clotting method according to Clauss.

- 10 -

## Example 2

The Effect of Various Concentrations of Tranexamic Acid and Arginine-HCl on the force/mm (slope) of the clot

	Group A	Group B	Group C	Group D	Group E	Group F	Group G
	0% TEA	0% TEA	5% TEA	10% TEA	10% TEA	10% TEA	10% TEA
	0% Arg.	2% Arg.	2% Arg.	2% Arg.	0% Arg.	1% Arg.	4% Arg.
incubation days							
0	2.34	3.55	3.49	3.1	3.92	3.26	0.68
1	0.9	1.9	3.3	3.17	3.4	3.2	1.83
4	0.5	1.13	3.24	3.02	3.49	3.17	1.3
7			2.95	2.79	3.49	3.12	1.75
9			2.92	2.71	3.37	3.21	1.54
11			2.94	2.73	3.34	3.19	1.16
14			2.86	2.82	3.37	3.08	

Conclusion: At freezing either Arg. or TEA enhance the slope of the clot (the clot is stronger than without stabilizers but TEA has stabilizing effect in the strength of the clot with time when incubating at 4 - 8°C.

- 11 -

Example 3  
The Effect of Various Concentrations of Tranexamic Acid and Arginine-HCl on the Content of Clottable Fibrinogen

	Group A	Group B	Group C	Group D	Group E	Group F	Group G
	0% TEA	0% TEA	5% TEA	10% TEA	10% TEA	10% TEA	10% TEA
	0% Arg.	2% Arg.	2% Arg.	2% Arg.	0% Arg.	1% Arg.	4% Arg.
storage at - 70°C	31.96	37.22	41.18	40.91	41.79	42.21	39.46
incubation days							
0	35.2	34.01	40.92	45.46	40.99	43.61	41.65
1	26.23	32.77	41.38	41.61	39.82	40.47	39.2
4	22.5	21.41	42.03	41.02	37.08	39.56	40.46
7	24.7	18.07	37.98	42.25	41.04	40.04	38.5
9	23.37	18.4	39.61	40.47	39.28	37.02	37.61
11	20.16	12.58	37.82	39.39	37.73	39.46	37.28
14	23.91	15.31	39.32	39.38	40.49	38.43	40.79
30	22.13	13.24	35.24	39.13	40.47	39.53	37.15

Conclusion: TEA stabilizes clottable fibrinogen

Example 4  
The Effect of Various Concentrations of Tranexamic Acid and Arginine-HCl on Factor VIII Activity

	Group A	Group B	Group C	Group D	Group E	Group F	Group G
	0% TEA	0% TEA	5% TEA	10% TEA	10% TEA	10% TEA	10% TEA
	0% Arg.	2% Arg.	2% Arg.	2% Arg.	0% Arg.	1% Arg.	4% Arg.
storage at - 70°C	9.13	11.15	13.98	15.26	12.87	13.51	14.23
incubation days							
0	8.77	9.19	11.22	11.87	8.42	10.52	10.67
1	6.54	3.38	9.79	10.15	8.91	8.11	6.85
4	4.78	6.32	9.16	10.28	8.68	8.93	9.15
7	6.19	5.61	8.98	11.31	8.84	11.6	11.73
9	3.62	2.48	7.77	10.19	8.69	8.12	7.23
11	4.27	3.13	8.55	10.16	9.32	9.17	9.31
14	5.61	3.37	8.89	10.13	8.66	8.09	8.46

Conclusion: Arg. stabilizes FVIII at freezing and TEA stabilizes FVIII during incubation at 4 - 8°C.

C l a i m s

1. A mixture comprising fibrinogen, 4-(aminomethyl)-cyclohexane-carboxylic acid (tranexamic acid) as well as arginine, lysine or combinations thereof.
2. The mixture of claim 1 comprising a solution of fibrinogen, tranexamic acid as well as arginine, lysine and/or combinations thereof.
3. The solution of claim 1 and/or 2 comprising a biological active component (BAC) which is a solution of proteins derived from blood plasma comprising fibrinogen, tranexamic acid and arginine or lysine or mixtures or arginine and lysine, their pharmaceutically acceptable salts, as well as, optionally, substances forming a buffered solution in an aqueous medium.
4. The solution of any one of the claims 1 - 3, wherein the amount of tranexamic acid is from about 1 % - 20 % by weight and the amount of arginine or lysine hydrochloride is from about 0.1 % - 4 % by weight.
5. The solution of any one of the claims 1 - 3, wherein the amount of tranexamic acid is about 5 % - 15 % by weight and the amount of arginine or lysine hydrochloride is about 1 - 3 % by weight.
6. The solution of claim 3, wherein BAC is derived from a concentrated cryoprecipitate.
7. The solution of any one of the claims 1 to 6, wherein the BAC comprises a fibrinogen content of from about 15 to 150 mg/ml.
8. The solution of anyone of the claims 1 to 7, wherein the BAC comprises further factor VIII, factor XIII,

- 14 -

fibronectin, von Willebrand factor (vWF), vitronectin.

9. The solution of claim 8, wherein the factors VIII, XIII, fibronectin, von Willebrand factor (vWF), vitronectin have been prepared by recombinant methods.
10. The solution of any one of the claims 1 to 9, wherein the buffered solution is a glycine buffer.
11. A two component tissue glue comprising separately the components A and B, wherein
  - component A comprises a solution of one of the claims 1 to 10, and
  - component B comprises a solution of a proteolytic enzyme which is capable of forming fibrin when it reacts with fibrinogen.
12. The two component tissue glue of claim 11, wherein the component B comprises human thrombin in an activity of from about 2 - 4000 international units per ml measured according to the method of Clauss.
13. The two component tissue glue of any one of claim 11 or 12, wherein component B is stabilized with tranexamic acid and arginine.
14. Fibrin clot obtainable by mixing component A and component B in an appropriate ratio, preferably 1 : 1.



- 1 / 1 -




FIG.1

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 98/00502

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61L25/00 A61K38/36 C07K14/75

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61L A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 22503 A (OPPERBAS HOLDING B V ; MARTINOWITZ URI (IL); BAL FREDERIC (AT)) 13 October 1994 cited in the application see abstract see page 4, paragraph 2 see page 6, paragraph 3 see page 7, paragraph 2 see claims 1,2,8 ---	1-3,6,11
X	US 5 578 326 A (WEIS-FOGH ULLA S) 26 November 1996 see abstract see column 7, line 51 - column 8, line 60 --- -/--	1-3

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

27 May 1998

Date of mailing of the international search report

08/06/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx: 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Heck, G

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 98/00502

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TAKADA, A. ET AL: "Importance of lysine in the regulation of plasma proteases and inhibitors" ANN. UNIV. SARAV.: MED., SUPPL. (1983), 3(ANTITHROMBIN 3: BIOCHEM., FUNCT., ASSAY, CLIN. SIGNIF.), 17-22 CODEN: AUSSEI;ISSN: 0173-6973, 1983, XP002034352 see page 19, paragraph 1-3; figure 7 see page 20, paragraph 1; figure 8 ---	1-5,7, 11-14
T	LENTNER C.: "GEIGY SCIENTIFIC TABLES" 1981, CIBA-GEIGY, BASEL XP002034353 212490 see page 92 - page 93; tables 2-4 -----	1-3,6,11

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/00502

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9422503 A	13-10-1994	AT 151295 T	15-04-1997
		AU 4566893 A	14-02-1994
		AU 6536394 A	24-10-1994
		CA 2140427 A	03-02-1994
		CA 2159469 A	13-10-1994
		CZ 9500054 A	14-06-1995
		DE 69309685 D	15-05-1997
		DE 69309685 T	16-10-1997
		WO 9402182 A	03-02-1994
		EP 0651659 A	10-05-1995
		EP 0691858 A	17-01-1996
		FI 950194 A	17-01-1995
		HU 71873 A	28-02-1996
		JP 8502035 T	05-03-1996
		JP 8512215 T	24-12-1996
		NO 950171 A	17-01-1995
		SK 5095 A	11-07-1995
US 5578326 A	26-11-1996	AU 8076687 A	21-04-1988
		CA 1320439 A	20-07-1993
		DE 3777398 A	16-04-1992
		WO 8802259 A	07-04-1988
		EP 0287609 A	26-10-1988
		US 5318782 A	07-06-1994